

PQ 10b Q

Q1

A 3 kg mass is moving at 6 m s^{-1} .

- (a) Calculate its kinetic energy (E_k).
- (b) Calculate the work required to change its speed to 12 m s^{-1} .
- (c) If the work is done by a force of 2.5 N, calculate how far the mass moves while the force is acting.
- (d) How long does this take?
- (e) What power is developed by the force?

Q2

A 0.25 kg toy car has 8 joules of E_K .

- (a) Calculate its speed.
- (b) How much work is required to be done to double the speed of the car?
- (c) What force is required to stop the car over a distance of 6.4 m?
- (d) How long will this take?
- (e) What power is developed by the force?

Q3

A 1.4 kg rock is fired from ground level straight up into the air at 5 m s^{-1} . Find:

- (a) Its initial kinetic energy.
- (b) How high above the ground it rises.
- (c) Its gravitational potential energy at its highest point.
- (d) The work gravity will do in returning the ball to the ground.

Q4

An object moving at 3 m s^{-1} has 54 J of E_k .

- (a) Calculate its mass.
- (b) What work is done to stop this object?
- (c) If the work is done by a constant frictional force of 2N , how far will it move while stopping?
- (d) Find the acceleration of the object.
- (e) How long will it take to stop?
- (f) What power will the force develop in stopping the object?

Q5

Calculate the work required to stop each of the following moving objects.

- (a) A 40 kg mass moving at 0.1 m s^{-1} .
- (b) A 0.2 kg mass moving at 10 m s^{-1} .
- (c) A 2 kg mass moving at 5 m s^{-1} .
- (d) A 4.0 kg mass moving at 7 m s^{-1} .
- (e) A 0.1 kg mass moving at 20 m s^{-1} .
- (f) Each object is stopped by a force of 6 N. How far does each object move while stopping?

Q6

A 2 kg object, moving at 5 m s^{-1} accelerates until it reaches 15 m s^{-1} . This takes 4.0 s. Calculate:

- (a) Its acceleration.
- (b) The kinetic energy it gained.
- (c) The work done on the object.
- (d) How fast it would be moving if its E_k doubled from its value at 15 m s^{-1} .
- (e) The extra work needed to be done to increase its speed from 15 m s^{-1} to 50 m s^{-1} .

Q7

Calculate by how much the kinetic energy of an object changes if:

- (a) Its mass doubles.
- (b) Its mass triples.
- (c) Its mass quadruples.
- (d) Its speed doubles.
- (e) Its speed triples.
- (f) Its speed quadruples.

Q8

An engine develops 20 000 W of power in moving a 250 kg car against a frictional force of 500 N. This takes 4 s and the car moved a distance of 30 m. Find:

- (a) The total work done.
- (b) The work done against friction.
- (c) The increase in the E_K of the car.
- (d) The car's speed after the 4 s.

Q9

A 40 kg model car accelerates from rest to 20 m s^{-1} against a frictional force of 0.25 N kg^{-1} . This takes 8 s. Find:

- (a) The acceleration of the car.
- (b) The force causing the acceleration.
- (c) The total force acting on the car.
- (c) The work done on the car.
- (d) How far the car travels.
- (e) The work done against friction.
- (f) The total work done by the force.
- (g) The power developed by the car.